

SPECIES COMPOSITION OF PHYTOPLANKTON IN THE TWO MIGRATORY BIRDS VISITED PONDS OF KANYAKUMARI DISTRICT

G. S. REGINI BALASINGH¹, S. JEEVA² & W. VINCY³

Department of Botany and Research Centre, Scott Christian College (Autonomous),
Nagercoil, Tamil Nadu, India

ABSTRACT

The present investigation was carried out from January 2012 to December in order to determine the seasonal variation and species composition of phytoplankton in the two migratory birds visited ponds of this district. Samples were collected from the two pond for the analysis of physico-chemical parameters and phytoplankton. The dissolved oxygen concentration ranged for a 4.8ml/l to 6.18mg/l, Nitrate 1.03 to 1.45mg/l, Phosphate 0.12 to 0.18mg/l. A total of 39 genera, 55 species in pond A and 45 genera 55 species in pond B were recorded under 4 groups viz. Chlorophyceae, Cyanophyceae, Bacillariophyceae and Euglenophyceae with maximum diversity of Cyanophyta.

KEYWORDS: Phytoplankton, Physico-Chemical Parameter, Pollution, Diversity, Cyanophyta

INTRODUCTION

Phytoplankton as primary producer plays a pivotal role in the aquatic ecosystem as most of the solar energy is changed into food energy by these organisms. Phytoplankton includes several thousands of microalgae belonged to chlorophyta (green algae), cyanophyta (blue green algae), Bacillariophyta (diatoms), dinoflagellates, cryptomonads etc. They respond quickly to environmental changes and are used to assess the ecological status of water quality. Several algal species are used as pollution indicators, (Palmer, 1969). The distribution and dynamics of algal species varies depending upon the physico-chemical parameters like temperature, light intensity, biological oxygen demand, dissolved oxygen, alkalinity and concentrations of micro and macro elements, (Tiwari *et al.*, 2004).

Several investigations were carried out in the phytoplankton diversity and their relationship with abiotic factors, (Pahari and Dutt 2000; Perumalsamy and Thangamony 2004; Jena *et al.*, 2006; Tiwari and Chauhan 2006b; Malliswar *et al.*, 2007; Panikar *et al.*, 2012). In kanyakumari district the migratory birds visited few freshwater environments and no previous detailed ecological study regarding the phytoplankton diversity in this regard. Hence the present investigation was carried out (2012 January to December) to determine the species composition of phytoplankton in two different water bodies of migratory visited ponds of the district.

MATERIALS AND METHODS

Phytoplankton samples were collected from the two freshwater ponds (Pond A- Parakai kulam; Pond B- Puthari kulam) on the monthly basis from January 2012 to December 2012 by using 1 litre sterilized wide mouth plastic containers at each pond (APHA, 1985) and fixed in 4% formalin solution. Identification and enumeration was carried out by relevant literatures (Round 1971; Anand, 1998; Prescott 1978 and Krishnamurthy, 2000).

RESULTS

Physico-chemical parameters of water is the prime consideration to assess its quality for irrigation, fishing, boating and industrial purposes. The phytoplankton abundance and succession linked with the influence of abiotic factors. In Pond-A and Pond-B a total of 39 genera 55 species and 45 genera 69 species phytoplankton members were reported and documented in Table 2 and they belonged to 4 classes viz.. Cyanophyceae, Chlorophyceae, Bacillariophyceae and Euglenophyceae. In pond-A, 16 genera 24 species to cyanophyta, 11 genus with 14 species to chlorophyta, 9 genera, 13 species to Bacillariophyta, 3 genera with 4 species to Euglenophyta were recorded. In the second pond (Pond-B) 18 genera 31 species to Cyanophyta, 17 genera 24 species to Chlorophyta, 7 genera 8 species to Bacillariophyta, 3 genus with 6 species to Euglenophyta were reported. Cyanophycean members contributed maximum diversity in both the ponds. Genus like *Chroococcus*, *Microcystis*, *Oscillatoria*, *Merismopedia*, *Anabaena*, *Nostoc* and *Aphanocapsa* were found to be dominant while *Eucapsis*, *Coelosporium*, *Rivularia*, *Arthrospira* and *Synechocystis* were reported abundant. *Phormidium*, *Anacystis*, *Lyngbya* and *Aphanothece* were just recorded.

In the case of Chlorophyceae *Ankistrodesmus*, *Chlorococcum*, *Pediastrum*, *Secnedesmus*, *Closterium*, *Spirogyra*, *Crucigenia*, *Tetraspora* were dominant. *Oedogonium*, *Ulothrix*, *Cladophora*, *Botriyococcas* and *Chlorella* were found to be abundant while *Netrium*, *Tetradon* and *Hydrodictyon* were recorded occasionally.

Among the diatoms *Navicula*, *Pinnularia*, *Fragillaria*, *Synedra* and *Gamponema* were formed to be the dominant genera while *Amphora*, and *Achnanthes* were abundant. *Diatoma* and *Surirella* were recorded rarely.

As for as Euglenophyceae is concerned the genus *Euglena* and *Phacus* were recorded at certain areas of the pond.

During the study period phytoplankton cell density reached maximum in April (14.48×10^3 cells/ m³) and October (12.87×10^3 cells/m³) showing a bimodal peak in Pond-A and a unimodel peak was noticed during December (16.9×10^3 cells/m³) in the second pond-B.

The physico-chemical parameters of the study pond showed higher mean p^H (7.74 ± 0.51), temperature (27.29 ± 2.07) along with higher dissolved oxygen concentrations (6.18 ± 0.38 mg/l). The levels of potassium was maximum (7.42 mg/l) in Pond-A while nitrate and phosphate content remains maximum in Pond-B and minimum in Pond-A (Table.1).

DISCUSSIONS

Phytoplankton are sensitive to the environmental changes and their distribution varies considerably with respect to seasons, water quality and nutrient concentrations (Thirugnamoorthy and Selvaraju, 2009; Ganai *et al.*, 2010b). Its biomass in a particular time is a dynamic process altered by the interaction of light, nutrient availability, growth and reproduction of plankton and cropping by herbivores (Diana *et al.*, 1997). Similar to the present investigation Devi *et al* (2007), Rout and Borah (2009), Mahar and Beena (2010) have reported four algal groups, viz. Chlorophyta, Cyanophyta, Bacillariophyta and Euglenophyta in their studies. Among the four groups, Cyanophyceae members dominated and earlier observation of Hudder (1995), Kavitha (2006), Hujare (2008b) and Joseph (2012) also pointed out the same in their studies. The cell density of phytoplankton reached maximum during monsoon or non-monsoon or both the months exhibiting uni or bimodal peaks. Pond-A represent bimodal peaks in summer and northeast monsoon season whereas Pond-B showed unimodel peak during northeast monsoon season. Similar findings of unimodel peak was reported by Bhagat *et al.*, (2009)

and Ganai *et al.*, (2010b). Srivastava (2002) and Chellapa *et al.*, (2009) has also pointed out the existence of bimodal peaks of phytoplankton. Many workers have reported blue green algae as a dominant group during non-monsoon season (Zacharias and Roy 2007; Hussan *et al.*, 2008; Khanna and Indu, 2009) and north east monsoon season (Santhosh *et al.*, 2007 and Jeyabaye, 2010). The dominant nature of Cyanophycean members are the characteristic feature of eutrophic environment which have high concentrations of nutrient especially phosphate and nitrate (Neelam *et al.*, 2009). The waste excreta, dead remnants of birds and fishes has increased the organic content of the pond and hence the number of cyanophycean members reached maximum in their density. Similar trend was reported in koonthankulam bird sanctuary pond by Esaki (2006). Moreover a direct effect of temperature on algal growth may influence the growth of phytoplankton when the temperature was above 23°C to 30°C it favours the growth of Cyanophyceae. In the present observation maximum population of algae was during summer months when the temperature was between 25-30°C. The p^H range between 6.7 to 7.8 (Healey, 1982) alkalinity range of 50 to 110mg/l (Jackson, 1971), high nutrient concentration (Ghavzan and Gunale, 2006 and Neelam *et al.*, 2009), rich organic matter (Kant and Amita, 1999) and dissolved oxygen (Ansari *et al.*, 2008) of the two pond produces maximum diversity of cyanophycean members. The genus *Microcystis aeruginosa* produced blooms during November month in Pond-A while *Chroococcus turgidus* produced blooms during December in the second pond.

The second dominant group in both the pond was Chlorophyta with a total of 43 and 25 species respectively. Among the Chlorophycean members the genus *Pediastrum* was reported with four species, *Scendesmus* and *Closterium* with three species each. Similar findings were made by Ida *et al.*, (2004) and Shamel (2012) in their studies on phytoplankton diversity. The third group reported was Bacillariophyta with 11 genera and the genus *Navicula* includes four species Euglenophyta was represented with three genus which indicated the presence of polluted nature of ponds (Balasingh and Shamel, 2007; Desh *et al.*, 2010). Previous studies on freshwater environment showed that higher temperature and nitrate concentration favours the growth of euglenoids in the two experimental ponds (Jasprica *et al.*, 2006). Blue green algae such as *Oscillatoria*, *Microcystis*, *Chroococcus* and Euglenophycean members like *Phacus* and *Trachelomonas* were reported as pollution indicators (Palmer, 1969). Devi and Charya (2007) observed that the members like *Microcystis* and *Lymbya* were the indicators of bad water quality.

CONCLUSIONS

From this study it can be concluded that the rich organic waste of the freshwater environment, enriched by the excreta, has increased maximum Cyanophycean diversity.

ACKNOWLEDGEMENTS

The author is thankful to UGC for all financial support and helps rendered by the department of Botany. I thank the Principal, Scott Christian College for all moral support.

Table 1: Annual Mean Values of Physico-Chemical Parameters in Pond-A and Pond-B during the Year 2012

Parameters	Pond-A	Pond-B
Temperature	27.29 ± 2.07	24.97 ± 2.68
p ^H	7.74 ± 0.51	7.64 ± 0.38
D.O (mg/l)	6.18 ± 0.38	4.8 ± 1.01
BOD (mg/l)	2.26 ± 0.42	2.6 ± 0.74
Total alkalinity (mg/l)	62.14 ± 11.08	53.78 ± 8.17

Table 1 – Cond.,		
Calcium (mg/l)	12.23 ± 2.40	11.38 ± 1.68
Potassium (mg/l)	7.42 ± 2.19	6.18 ± 2.40
Nitrate (mg/l)	1.03 ± 0.28	1.45 ± 0.54
Phosphate (mg/l)	0.18 ± 0.03	0.12 ± 0.02

Table 2: Phytoplankton Species Recorded in the Ponds-A and Pond-B during the Year 2012

S. No	Name of the Algae	Pond-A	Pond-B
CYANOPHYTA			
1.	Aphanocapsa biformis A.Br.	+	+
2.	Aphanothece sp	+	+
3.	Chroococcus tenax (Kirchn) Hieron	-	+
4.	Chroococcus indicus, Zeller.	+	-
5.	Chroococcus minor (Kuetz) Nag	+	+
6.	Chroococcus turgidus (Kuetz) Nag	+	+
7.	Chroococcus minutus (Kuetz)Nag.	+	+
8.	Coelosphaerium sps.	+	+
9.	Eucapsis minuta Fritsch	+	+
10.	Gomphosphaeria aponina Kuetz.	+	+
11.	Merismopedia punctata Meyen	-	+
12.	Merismopedia minima Beck.	+	+
13.	Merismopedia elegansLemm	-	+
14.	Microcystis aeruginosa Kuetz.	+	+
15.	Microcystis protocystis Crow	-	+
16.	Microcystis flos-aquae (Wilttra) Kirchner	-	+
17.	Microcystis elongata Desikachary	+	+
18.	Synechococcus elongatus Nag	+	+
19.	Arthrospira platensis (Nordstedt) Gomont	+	+
20.	Lyngbya spiralis Geitler	+	+
21.	Oscillatoria formosa Bory ex Gomont	+	+
22.	Oscillatoria limosa Ag. Ex Gomont	-	+
23.	Oscillatoria princeps Vaucher ex Gomont	-	+
24.	Oscillatoria tenuis Ag. ex Gomont	+	+
25.	Oscillatoria sancta (Kuetzing) Gomont	+	+
26.	Phormidium fragile (Meneghini) Gomont	+	+
27.	Anabaenopsis circularis (G.S.West)Wolosz. et Miller	+	+
28.	Anabaena orientalis Dixit.	+	+
29.	Nostoc commune Vaucher ex Born. et Flah.	+	+
30.	Nostoc muscorum Ag.	-	+
31.	Anacystis sps	+	+
32.	Rivularia aquatica De Wilde.	+	+
CHLOROPHYTA			
33.	Ankistrodesmus falcatusvar.acicularis(Corda.)	-	+
34.	Ankistrodesmus convolutesCorda.	+	+
35.	Chlorococcum humicola (Nag.) Raban.	+	+
36.	Chlorella vulgarisBayernick	+	+
37.	Coelastrum indicum Turner.	+	+
38.	Hydrodictyon sps	-	+
39.	Crucigenia tetrapedia(Kirchner)	+	+
40.	Pediastrum angulosum (Ehr.) Menegh.	+	+
41.	Pediastrum duplex Lagerheim	+	+

42.	<i>Pediastrum integrum</i> Nagelli	+	+
43.	<i>Pediastrum tetras</i> (Ehr.) Ralfs.	-	+
44.	<i>Botriyococcus</i> sps	-	+
45.	<i>Scenedesmus armatus</i> var. <i>bicaudatus</i> Smith.	-	+
46.	<i>Scenedesmus bijugatus</i> var. <i>alternans</i> Kuetz.	-	+
47.	<i>Scenedesmus quadricauda</i> var. <i>quadrispinia</i> Breb.	+	+
48.	<i>Tetraedron gracile</i> (Rein.) Hansgirg	-	+
49.	<i>Tetradporu</i> Sps	-	+
50.	<i>Ulothrix zonata</i> (Kuetz)	+	-
51.	<i>Oedogonium hispidum</i> Nordstedt	+	+
52.	<i>Closterium acerosum</i> (Schränk.) Ehr.	-	+
53.	<i>Closterium ehrenbergii</i> (Menegh.) ex Ralfs.	+	+
54.	<i>Cosmarium turgidum</i> Ralfs	+	+
55.	<i>Netrium digitus</i> (Ehrenberg) Roth.	+	+
56.	<i>Pithophora</i> sps	-	+
57.	<i>Spirogyra gratiana</i> Transean.	-	+
BACILLARIOPHYTA			
58.	<i>Amphora ovalis</i> Kuetz	+	-
59.	<i>Cymbella lanceolata</i> Breb.	-	+
60.	<i>Fragilaria intermedia</i> (Grun.)	+	+
61.	<i>Gomphonema lanceolatum</i> Her.	+	+
62.	<i>Navicula cuspidate</i> Kuetz	+	+
63.	<i>Navicula radiosa</i> Kuetz.	+	-
64.	<i>Navicula halophilakuetz.</i>	+	-
65.	<i>Navicula viridula</i> Kuetz	+	+
66.	<i>Achnanthes inflata</i>	+	-
67.	<i>Pinnularia simplex</i> Ehr.	+	-
68.	<i>Pinnularia viridis</i> (Nitzsch.) Ehr.	+	+
69.	<i>Synedra ulna</i> (Nitzsch.) Ehr.	+	-
70.	<i>Diatoma</i> Sps	+	-
71.	<i>Suriella elegans</i> Ehr.	-	+
72.	<i>Tabellaria flocculosa</i> (Roth) Kuetz.	+	+
EUGLENOPHYTA			
73.	<i>Euglena gracilis</i> Mallisch	-	+
74.	<i>Euglena sanguine</i> Ehrenberg	+	+
75.	<i>Phacus aribicularis</i> Var. <i>Caudata</i>	+	+
76.	<i>Phacus cylindricus</i> Pochmann	+	+
77.	<i>Phacus curvicauda</i> Swir. Skz.	+	+
78.	<i>Trachelomonas volvocina</i> Ehr.	-	+

REFERENCES

1. Anand, N., 1998. Indian freshwater Microalgae Bishen Singh Manendra-Palsingh, Dehradun, 1. Anand India, pp. 5-88.
2. Ansari, M.F., Ankalg, R.E. and S.R. Ankalg, 2008. Studies on Physico-chemical and plankton of Unkal lake at Hubi (Karnataka, India). In Sengupta M. and R. Dalwani (eds) Proc. Of Taal 2007. The 12th Lake World Conference, pp. 1687-1694.
3. APHA, 1985. Standard methods for the Examination of Water and Waste Water Washington, D.C. USA.

4. Balasingh, G.S.R., and V.P. Selva Shamal, 2007. Phytoplankton diversity of a perennial pond in Kanyakumari District. *Journal of Basic and Applied Biology*, 1 (1): 23-26.
5. Bhagat, P., Ragini Gupta and P.K. Gupta, 2009. Seasonal abundance of phytoplankton community in Lake Naukuchiyatal, Uttarakhand, India. *J. Ecophysiol. Occup. Hith.*, 9: 171: 182.
6. Chellapa, N.T., Camara, F.R.A and Rocha, O. b., 2009. Phytoplankton community Indicators of water quality in the Armando- Ribeiro Goncalves Reservoir and Patayo channel Rio Grande do Norte, Braz. *J. Biol.* 69(2): 241-251.
7. Desh, P.K., Mohapatra, P.K. and M. Kar, 2010. Freshwater algal diversity of Northern Eastern Ghats, India. *IEPTRI ENVIS News letter*, 16 (1): 3 – 6.
8. Devi Anita, U., and Charya, M.A.S, 2007. Phytoplankton in lower Manair dam and kakatica canal, Karimnager, Nat. Environ. Pollu. Tech. 6(4): 643-648.
9. Diana, J.S., Szyper, J.P., Batterson, T.R., Boyd, C.E., and Piedrahita, R.H., 1997. Water quality in pond. In: Dynamics of Pond Aquaculture.(ed. By H.S.Egna & C.E.Boyd) CRC Press. New York, pp: 33.
10. Esakki, G., 2006. Ecophycological investigation of Koondhankulam bird sanctuary, Tirunelveli District. M.Phil. Dissertation, Manonmaniam Sundaranar University.
11. Ganai, A.H., Parveen, S. and Asif A. Khan, and Maryam, H., 2010b. Phytoplankton diversity at Watlab Ghat in Wular lake, Kashmir. *J. Eco. Nat. Environ.* 2(8): 140-146.
12. Ghavzan Jafari, N., and Gunale, V.R., 2006. Studies on correlation between algae and physico-chemical characteristic from mula river, Pune, India. *Eco. Environ. and Cons.* 12(2): 369-376.
13. Healey, F.P., 1982. Phosphate In. Carr. N and B. Whitton (eds). The biology of Cyanobacteria. Blackwell Scientific Publishers. Pp. 105- 107.
14. Hudder, B.D., 1995. Hydrobiological studies in lentic freshwater bodies of Hubli Ph.D. Thesis. Submitted to Karnataka University, Dharwad, pp. 267.
15. Hujare Milind, S., 2008b. Seasonal variation of phytoplankton in the freshwater tank of Talsande, Maharastra. *Nat. Envir. And Pollu. Tech.* 7(1): 147-150.
16. Ida, C.J., Palavesam, A., G. Santhanakumar and A.D.S. Raj, 2004. Dynamics of phytoplankton in relation to limnological conditions of a perennial pond in Kanyakumari district. *Indian Journal of Theoretical and Experimental Biology*, 1: 35-42.
17. Jackson, M.L., 1973. Soil Chemical Analysis Prentice Hall of India Pvt. Ltd. New Delhi.
18. Jasprica, N., Dubravika, H., Marina, C., and Rimac, A., 2006. A preliminary investigation of phytoplankton of karstic pond (Dugi otok Island, Croatia) Acta. Bot. Croat., 65(2): 181-190.
19. Jayabhaye, U.M., 2010. Studies on Phytoplankton diversity in Sawana Dam, Maharastra, India. Samiksha Aur Mulyankan., 11(11-12): 40-42.

- Jena, M., S.K. Radha and S.P. Adhikary, 2006a. Algal diversity of similipal biosphere reserve, Orissa. *Indian Hydrobiology*, 9(1): 103-113.
20. Joseph, J., 2012. Studies on the cyanobacterial diversity from selected freshwater environment and its application in biotechnology. Ph.D. Thesis. Submitted to M.S. University, Tirunelveli.
21. Kant, S., and Amita, S., 1999. Algal diversity of Banganga (Katra). *Proc. Acad. Environ. Biol.*, 8(1): 15-24.
22. Kavitha, A., 2006. Ecotechnological application of Cyanobacteria for microproductivity enhancement in fresh water agricultural system. Ph.D. Thesis, Manonmaniam Sundaranar University, Tirunelveli, Tamilnadu.
23. Khanna, V., and Indu. Y., 2009. Planktonic diversity in the holy lake of Pushkar Ajmar. *Nat. Envir. Pollu. Tech.*, 8(2): 339-342.
24. Krishnamurthy, V., 2000. Algae of India and neighbouring countries I. Chlorophyta, Oxford and IBH Publishing Co. Pvt. Ltd New Delhi. Pp.210.
25. Malliswar, V.N.S., Krupanidhi, D.S., and Chandra, S., 2007. Algal diversity in a man made water body, *Indi. Hydro*. 10(1): 79-86.
26. Mohar, R.K and S.Beena, 2010. Diversity and seasonal fluctuation of phytoplankton in freshwater reservoir Ighira, Gualior. *Inter. Res. Jour.* 2010.
27. Neelam, V., Avinash, B., and Dwivedi, S.N., 2009. Planktonic Biodiversity of Bhoj. Wetland, Bhopal, India. *J. Appl. Sci. Environ. Manage.*, 13(4): 103-111.
28. Pahari, P.R., and Dutt, T.K., 2000. Macrophyte Planktonic Components of two contrasting wetlands in Midnapore district. *Env. Eco.* 18(1): 196-201.
29. Palmer, C.M., 1969. A composite rating of algae tolerating organic pollution. *J. Phycol.*, 5: 78-82.
30. Panikkar, M.V.N., Jayalakshmi, R. and Jackson, A. 2012. Species diversity of two chlorophycean colonial genera (*Pediastrum meyen* and *Scenedesmus meyer*) from the freshwater ecosystem of kerala. *Ind. Hydro*, 14(2): 109-116.
31. Perumalsamy, K. and Thangamani, 2004. Indices of zooplankton richness and evenness and bioindicators in some perennial ponds, Koassay, 17(12): 35-39.
32. Prescott, C.W., 1978. Algae of Western great lakes area. Wm C. Brown Co. Publishers. Dubuque Iowa, pp. 977.
33. Prescott, G.W., 1978. How to know freshwater algae 3^{ed}. Wes, C. Brown Company Publication, Jowa, USA. 1-280pp.
34. Round, F.E. 1971. The taxonomy of the Chlorophyta 11. *Br. Phycol. J.* 6: 235 64.
35. Rout J. and D. Borah, 2009. Algal diversity in Chatla wetland in Cachar District (Southern Assam). *Assam Uni. J. of Sci and Technology: Biol., Sci.*, 4(1): 46 – 55.
36. Santhosh, S.V., Shoba, S., Girijalakshmi and Thara, J.C., 2007. Phytoplankton abundance of Paravarkappil backwater of kerala. *Seaweed. Res. Util.*, 29(112): 155-163.

37. Srivastava, M.L., 2002. Physico-chemical and microbial characters of water. Daya publishing house, New Delhi. Pp. 117-133.
38. Thirugana moorthy, K., and Selvaraju, M., 2009. Phytoplankton diversity in relation to physico-chemical parameters of Gnanaprakasam temple pond of Chidambaram in Tamil Nadu, India. *Recent Research in Science and Technology*, 1(5): 235-238.
39. Tiwari, A. and S.V.S. Chauhan, 2006b. Wall algae from Agra. *Ind. Bot. Soc.*, 80: 269-271.
40. Tiwari, S., Dixit, S. and Gupta, S.K., 2004. An evaluation of physico-chemical parameters in surface waters of Shahpura lake, Bhopal. *Poll. Res.* 25: 829-832.
41. Zacharias, R. and C.M. Joy, 2007. Algal biodiversity and succession in Periyar river at Aluva, Kerala. *Indian Hydrobiol.*, 10: 121-133.